

An upper limit to the masses of stars

Donald F. Figer

STScI, 3700 San Martin Drive, Baltimore, Maryland 21218, USA

There is no accepted upper mass limit for stars. Such a basic quantity eludes both theory and observation, because of an imperfect understanding of the star-formation process and because of incompleteness in surveying the Galaxy¹. The Arches cluster^{2–7} is ideal for investigating such limits, being large enough to expect stars at least as massive as ~ 500 solar masses ($\sim 500 M_{\odot}$; based on a typical mass function), and young enough for its most massive members to still be visible. It is also old enough to be free of its natal molecular cloud, it is at a well-established distance, and it is close enough for us to discern its individual stars². Here I report an absence of stars with initial masses greater than $130 M_{\odot}$ in the Arches cluster, whereas the typical mass function predicts 18. I conclude that this indicates a firm limit of $150 M_{\odot}$ for stars; the probability that the observations are consistent with there being no upper limit is 10^{-8} .

Theory provides little guide in determining the most massive star that can form. Pulsational instabilities were once thought to destroy stars more massive than $95 M_{\odot}$ (ref. 8); however, these pulsations may be damped⁹. Radiation pressure, and/or ionizing flux, inhibit accretion for stellar masses greater than $60 M_{\odot}$ (ref. 10), but direct collisions of protostellar clumps may overcome these effects¹¹. Although stellar evolution models have been computed for massive stars covering a large range in mass, up to $1,000 M_{\odot}$ (refs 12, 13), no such stars have ever been observed. Indeed, some of the most massive candidates have proved to be systems of multiple stars¹⁴.

Stars generally form with a frequency that decreases with increasing mass for masses greater than $\sim 1 M_{\odot}$, that is, $d(\log N)/d(\log m) = \Gamma$, where m is the initial stellar mass, N is the number of stars per logarithmic initial mass interval, and Γ is observed to be -1.35 (refs 15, 16). For stellar clusters young enough not to have lost members to supernovae, the distribution of stars is populated to the point where the mass function predicts one star, within the uncertainties of low number statistics. Therefore, stars with mass $M > 150 M_{\odot}$ can only be observed in very massive clusters with total stellar mass $> 10^4 M_{\odot}$. This requirement limits the potential sample of stellar clusters that can constrain the upper mass limit. Only a few clusters in the Galaxy satisfy this requirement, and all are located in the Galactic Centre.

To investigate the possibility that stars with $M > 150 M_{\odot}$ exist, imaging data were obtained using the Near-Infrared Camera and Multi-Object Spectrometer instrument on the Hubble Space Telescope in a programme to measure the mass functions of the most massive young clusters in the Galaxy, near the Galactic Centre². Intervening dust prevents observations of these clusters at optical or ultraviolet wavelengths, so images were obtained in near-infrared wavelengths (see Supplementary Fig. 1). Nearby control fields were also imaged to estimate the number of field stars that contaminate observations in such a densely populated region.

Photometry for stellar sources in the images was extracted, and corrected for the absorbing effects of dust by comparing the observed colours to those expected for the appropriate spectral types; note that intrinsic colours of massive stars on the main sequence at infrared wavelengths differ by only a few per cent. The dereddened fluxes were then converted into bolometric fluxes by accounting for the distance to the Galactic Centre, and the Geneva stellar evolution models were used to infer initial masses for each star¹⁷ (see Fig. 1). Although these models have associated errors,

note that the Arches stars are relatively unevolved; indeed, only the brightest dozen or so members show evidence of chemical enrichment by nucleosynthetic processes¹⁸. Some of the brightest stars in the cluster (three to ten, depending on cluster age within a range of 2–2.5 Myr and the coefficients in the extrapolation law) extend just above the $120 M_{\odot}$ limit of the mass–flux relation; I estimate masses for them that do not exceed $130 M_{\odot}$ through an extrapolation of this relation (see Supplementary Fig. 2).

The initial masses I estimate here agree with those inferred through wind/atmosphere modelling of high-resolution spectral observations to within a few per cent (ref. 3). Others have also applied the same technique to construct mass functions from infrared observations of massive young clusters, showing that these determinations are consistent with those estimated from optical observations¹⁹. In addition, several groups have found good consistency in physical properties inferred from optical and infrared analyses for massive stars at all stages of evolution^{20–22}.

Figure 2 shows the resultant initial mass function of the Arches cluster, assuming an age of 2 Myr, for stars within a projected radius of 0.5 pc, and solar metallicity^{2,18}. Although the cluster is the densest




Figure 1 Observed frequency distribution and inferred masses of stars in the Arches cluster versus brightness. Here $m_{F205W} = -2.5 \log(F/F_{Vega})$, where F is the flux of the star and F_{Vega} is the flux of Vega, as measured in the F205W ($\lambda_{\text{centre}} = 2.05 \mu\text{m}$) filter in NICMOS. **a**, Near-infrared luminosity functions of the central parsec of the Arches cluster (thin line) and nearby background fields (thick line). There are generally fewer bright than faint stars in both fields; but for the vast majority of the brightness range, there are more stars in the Arches cluster than in the control fields. This allows an accurate subtraction of background stars in order to create a mass function for the cluster. The shapes of the distributions are consistent with a very young stellar cluster in the cluster field and an old population ($\tau > 1$ Gyr) in the control fields. **b**, Inferred initial masses for Arches stars, based on the Geneva models¹⁷ for solar abundances and an age of 2 Myr. Each point represents one star in the cluster field. The three brightest stars have masses that slightly exceed $120 M_{\odot}$, the upper limit of the models, and are assigned masses through a linear extrapolation of the mass–flux relation from points immediately below this value.

in the Galaxy³, the data do not suffer from incompleteness due to crowding or sensitivity for the four highest mass bins in the figure. The small amount of background contamination was removed by subtracting the number of stars observed in nearby fields; this resulted in the subtraction of a total of seven stars from the upper four populated mass bins. The frequency distribution generally decreases with increasing mass, and is fitted by two lines through the four most massive populated bins, which contain 39 stars. One line has a slope of $\Gamma = -0.9$, appropriate for the most recent determinations^{2,23}, and the other has a slope equal to the Salpeter value that is observed for most clusters. For both slopes, there appears to be a deficit of expected very massive stars with masses beyond $\sim 130 M_{\odot}$; variations in assumed age (± 0.5 Myr), mass-loss rates and metallicity do not change the result. I estimate cumulative errors of $\sim 10\%$, and conclude conservatively that there is an upper mass cut-off of $\sim 150 M_{\odot}$ (see Supplementary Fig. 3 for the effects of mass loss on the most massive stars).

The observed deficit of stars is significant. If there is no upper mass cut-off, then the odds of identifying no stars beyond the observed limit are 10^{-8} if 18 are expected, and 10^{-14} if 33 are expected, assuming Poisson statistics. In addition, the maximum predicted stellar mass is at least $\sim 500\text{--}1,100 M_{\odot}$, values that are far beyond the masses inferred from the observations. I performed a Monte Carlo simulation of model systems to predict (as a function of cut-off mass) the probability that a cluster with the mass of the Arches cluster could have no stars with initial masses greater than $130 M_{\odot}$ (see Supplementary Fig. 4). In this simulation, I added uncertainties due to differential extinction, photometric error, average cluster age, a spread of ages for individual stars, and error in estimating the average cluster age. The simulation predicts few systems with no stars having initial masses greater than $130 M_{\odot}$ for cut-offs of $150 M_{\odot}$ or greater (Supplementary Fig. 4).

Clearly, the cluster age (τ) is an important quantity for the analysis. If the cluster is too old, $\tau > 3$ Myr, then its most massive

members would no longer be visible (that is, they would have progressed to supernovae), and the observations would then simply reveal an apparent cut-off due to the natural effects of stellar evolution. If the cluster is too young, $\tau \approx 1$ Myr, then the models would predict much higher initial masses for the brightest members; however, note that even younger ages would still require a firm upper mass cut-off, albeit at somewhat higher masses than predicted by the best estimated age. Analyses indicate that the cluster has an age of 2–2.5 Myr (refs 2, 3, 18). A younger age is inconsistent with the nitrogen-enriched atmospheres revealed in the spectra of the most massive stars in the cluster¹⁸. The fairly narrow age range is required by the observed heavy nitrogen enrichment in the brightest stars together with relatively weak observed nitrogen content in the atmospheres of slightly lower mass stars^{3,18}. An older age is inconsistent with the evolutionary status of the most massive stars in the cluster—they have not evolved to advanced stages, such as the carbon Wolf-Rayet phase^{3,6}. In addition, the lack of any supernova remnants in the cluster argues for an age less than 3 Myr. Indeed, if massive stars filling the apparent deficit were formed and evolved to supernovae, one would expect that a supernova remnant would have been formed at least every 50,000 yr for the past 0.5 Myr—yet none are observed. In summary, stars with masses above $\sim 150 M_{\odot}$ should still be visible if they were formed, given my estimate of the age for the Arches cluster.

The observed upper mass limit is on the low side of the estimated masses of a few massive stars in the Galaxy, although it still falls within the error bars of these estimates. It is important to note the large errors in such estimates. For instance, many of these estimates rely on stellar wind/atmosphere models that do not model the effects of increased opacity produced by metals in stellar winds (that is, line-blanketing). With more modern models, new mass estimates are smaller by up to a factor of two. In addition, mass estimates often suffer from uncertainties in distance, reddening and photometry. The typical build-up of errors can easily result in an uncertainty of a factor of two in flux, and of a similar factor in mass estimate. As an example, consider Pismis 24-1, which is estimated to have a mass of 210–290 M_{\odot} (ref. 24). The build-up in errors for this star, from effects described above, produces at least a factor of two variation in flux estimates, and the original mass estimates were produced without the use of line-blanketing. Once these combined effects are included, the true mass of this star may well be below 100 M_{\odot} . Note that uncertainty in distance is the next obstacle to making accurate mass estimates once line-blanketing is included; however, the distance to the Galactic Centre is very well known (to within 6%), and the Arches cluster is physically connected to phenomena known to be produced in the Galactic Centre³.

If there are stellar systems more massive than the limit, then perhaps they are binaries, or products of mergers of lower-mass stars. Indeed, the Pistol star, with an inferred initial mass of $\sim 150\text{--}250 M_{\odot}$ (ref. 13), is surrounded by Wolf-Rayet and red supergiant stars that are older than the expected lifetime of such a star²⁵. This apparent paradox may be reconciled if the star is actually multiple, or if it has recently experienced a rejuvenation through a merger with another star²⁶. High-spatial-resolution imaging suggests that the Pistol star is not binary to within a limit of 110 AU (ref. 13), yet massive binaries can have components with orbits on yet-smaller scales¹⁴.

An upper mass cut-off of $\sim 150 M_{\odot}$ was found for the cluster R136 in the low-metallicity environment of the nearby galaxy, the Large Magellanic Cloud²⁷. This result relies on an apparent deficit of ten stars with masses beyond this limit, based on the assumption that R136 has a total stellar mass of $5 \times 10^4 M_{\odot}$; however, this high cluster mass includes stars that span a range of ages, including those that exceed the age at which a massive star is expected to evolve to become a supernova. This has the effect of increasing the base of lower-mass stars from which to extrapolate an expected number of higher-mass stars, thus inflating an apparent deficit if those stars are




Figure 2 Frequency distribution versus mass for stars in the Arches cluster extracted from data in Fig. 1. The counts in each bin have been reduced by counts in nearby background fields. Error bars represent the Poisson errors based on the background subtracted counts. Two lines are drawn through the average counts in the four highest populated mass bins, with slopes inferred from the data ($d(\log N)/d(\log m) = \Gamma = -0.9$; ref. 2, 23) and that of Salpeter ($\Gamma = -1.35$; ref. 15). For both lines, there is a clear deficit of stars with initial masses greater than $\sim 150 M_{\odot}$, as seen in the hatched regions. In addition, both slopes predict that at least one star in the cluster should have a mass far beyond that observed if there is no upper mass cut-off. N_{missing} is the difference between the number of stars expected with $M_{\text{initial}} > 130 M_{\odot}$ and the number observed. M_{MAX} is the initial mass at which the mass function predicts the existence of one star. t_{age} is the assumed cluster age. Z is the assumed abundance of metals in the cluster stars.

not seen. Using a lower estimate of the cluster mass, $2 \times 10^4 M_{\odot}$ (ref. 28), in stars sufficiently young for the present analysis, I estimate that the true deficit beyond $150 M_{\odot}$ in R136 is roughly four stars—that is, the result in the present work is more statistically significant by this measure. If the deficit of massive stars in R136 is real, then it represents another measurement of the upper mass cut-off.

Surprisingly, the cut-off may be similar in environments that span a factor of three in metallicity^{18,29,30}, although metal content is often cited as a proxy for the source of opacity that limits the infall of material and eventual build-up of massive stars. This result implies that the process that limits the mass of a star is independent of metallicity, at least in the range of metallicities primarily found within the Galaxy and the nearby Large Magellanic Cloud. □

Received 5 October; accepted 9 December 2004; doi:10.1038/nature03293.

1. Elmegreen, B. G. Modeling a high-mass turn-down in the stellar initial mass function. *Astrophys. J.* **539**, 342–351 (2000).
2. Figer, D. F. *et al.* HST/NICMOS observations of massive stellar clusters near the Galactic center. *Astrophys. J.* **525**, 750–758 (1999).
3. Figer, D. F. *et al.* Massive stars in the Arches cluster. *Astrophys. J.* **581**, 258–275 (2002).
4. Nagata, T., Woodward, C. E., Shure, M. & Kobayashi, N. Object 17: Another cluster of emission-line stars near the Galactic center. *Astron. J.* **109**, 1676–1681 (1995).
5. Cotera, A. S. *et al.* A new cluster of hot stars near the Galactic center. *Bull. Am. Astron. Soc.* **24**, 1262 (1992).
6. Blum, R. D. *et al.* 2 micron narrowband adaptive optics imaging in the Arches cluster. *Astron. J.* **122**, 1875–1887 (2001).
7. Serabyn, E., Shupe, D. & Figer, D. F. An extraordinary cluster of massive stars near the centre of the Milky Way. *Nature* **394**, 448–451 (1998).
8. Schwarzschild, M. & Harm, R. On the maximum mass of stable stars. *Astrophys. J.* **129**, 637–646 (1959).
9. Beech, M. & Mitalas, R. Formation and evolution of massive stars. *Astrophys. J. Suppl.* **95**, 517–534 (1994).
10. Wolfire, M. G. & Cassinelli, J. P. Conditions for the formation of massive stars. *Astrophys. J.* **319**, 850–867 (1987).
11. Bonnell, I. A., Bate, M. R. & Zinnecker, H. On the formation of massive stars. *Mon. Not. R. Astron. Soc.* **298**, 93–102 (1998).
12. Bond, J. R., Arnett, W. D. & Carr, B. J. The evolution and fate of very massive objects. *Astrophys. J.* **280**, 825–847 (1984).
13. Figer, D. F. *et al.* The Pistol star. *Astrophys. J.* **506**, 384–404 (1998).
14. Daminieli, A. *et al.* η Carinae: binarity confirmed. *Astrophys. J.* **528**, L101–L104 (2000).
15. Salpeter, E. E. The luminosity function and stellar evolution. *Astrophys. J.* **121**, 161–167 (1955).
16. Kroupa, P. The initial mass function of stars: evidence for uniformity in variable systems. *Science* **295**, 82–91 (2002).
17. Schaller, G., Schaerer, D., Meynet, G. & Maeder, A. New grids of stellar models from 0.8 to 120 solar masses at $Z = 0.020$ and $Z = 0.001$. *Astron. Astrophys. Suppl.* **96**, 269–331 (1992).
18. Najarro, F., Figer, D. F., Hillier, D. J. & Kudritzki, R. P. Metallicity in the Galactic center: The Arches cluster. *Astrophys. J.* **611**, L105–L108 (2004).
19. Stolte, A., Brandner, W., Brandl, B., Zinnecker, H. & Grebel, E. K. The secrets of the nearest starburst cluster. I. Very Large Telescope/ISAAC photometry of NGC 3603. *Astron. J.* **128**, 765–786 (2004).
20. Bohannan, B. & Crowther, P. Quantitative near-infrared spectroscopy of OF and WNL stars. *Astrophys. J.* **511**, 374–388 (1999).
21. Hillier, D. J., Crowther, P. A., Najarro, F. & Fullerton, A. W. An optical and near-IR spectroscopic study of the extreme P Cygni-type supergiant HDE 316285. *Astron. Astrophys.* **340**, 483–496 (1998).
22. Najarro, F. Massive stars in the galactic center. *N. Astron. Rev.* **44**, 213–220 (2000).
23. Stolte, A., Grebel, E. K., Brandner, W. & Figer, D. F. The mass function of the Arches cluster from Gemini adaptive optics data. *Astron. Astrophys.* **394**, 459–478 (2002).
24. Walborn, N. R. *et al.* A new spectral classification system for the earliest O stars: Definition of type O2. *Astron. J.* **123**, 2754–2771 (2002).
25. Figer, D. F., McLean, I. S. & Morris, M. Massive stars in the Quintuplet cluster. *Astrophys. J.* **514**, 202–220 (1999).
26. Figer, D. F. & Kim, S. S. in *Stellar Collisions, Mergers and their Consequences* (ed. Shara, M.) 287–295 (ASP Conf. Ser. 263, Astronomical Society of the Pacific, San Francisco, 2002).
27. Weidner, C. & Kroupa, P. Evidence for a fundamental stellar upper mass limit from clustered star formation. *Mon. Not. R. Astron. Soc.* **348**, 187–191 (2003).
28. Walborn, N. R., Maiz-Apellaniz, J. & Barba, R. H. Further insights into the structure of 30 Doradus from the Hubble Space Telescope instruments. *Astron. J.* **124**, 1601–1624 (2002).
29. Ramirez, S. V. *et al.* Stellar iron abundances at the Galactic center. *Astrophys. J.* **537**, 205–220 (2000).
30. de Boer, K. S., Fitzpatrick, E. L. & Savage, B. D. Abundances of O, Mg, S, Cr, Mn, Ti, Ni and Zn from absorption lines of neutral gas in the Large Magellanic Cloud in front of R136. *Mon. Not. R. Astron. Soc.* **217**, 115–126 (1985).

Supplementary Information accompanies the paper on www.nature.com/nature.

Acknowledgements I acknowledge discussions with P. Najarro, R. Larson, N. Walborn, J. Puls, N. Panagia, M. Morris, C. Weidner, P. Kroupa, R. M. Rich, V. Bromm and M. Livio.

Competing interests statement The author declares that he has no competing financial interests.

Correspondence and requests for materials should be addressed to the author (figer@stsci.edu).